3 (Sem-3/CBCS) STA HC 3

2022 STATISTICS

(Honours)

Paper: STA-HC-3036

(Mathematical Analysis)

Full Marks: 60

Time: Three hours

The figures in the margin indicate full marks for the questions.

- 1. Answer **any seven** from the following questions: $1 \times 7 = 7$
 - (a) Identify the wrong statement:
 - (i) The set R of real numbers is the neighbourhood of each of its points.
 - (ii) The set Q of rationals is the neighbourhood of each of its points.
 - (iii) The open interval] a, b [is the neighbourhood of each of its points.

(Choose the correct option)

Contd.

- (b) The set $\left\{\frac{1}{n}:n\in\mathbb{N}\right\}$ has only one limit point, zero, which is not a member of the set. (State True or False)
- (c) If $\sum u_n$ is a positive term series, such that $\lim_{n\to\infty}\frac{u_n+1}{u_n}=l$, then the series converges if
 - (i) 1 < 1
 - (ii) l > 1
 - (iii) l=1
 - (iv) l=0 (Choose the correct option)
- (d) The positive term geometric series $1+r+r^2+....$ converges for r<1 and diverges for $r \ge 1$.

(State True or False)

- (e) Define alternating series.
- (f) A function which is continuous in a closed interval is also uniformly continuous in that interval.

(State True or False)

- (g) State the geometrical interpretation of Lagrange's mean value theorem.
- (h) State the expansion of $\cos x$.
- (i) The nth divided difference can be expressed as the product of multiple integrals. (State True or False)
- (j) Define the operators μ , δ used in calculus of finite differences.
- (k) State Stirling's formula for factorial n, when n is large.
- (1) Which of the following is not correct?
 - (i) Simpson's rule gives a better result than the trapezoidal rule.
 - (ii) Weddle's rule is generally more accurate than any of the others.
 - (iii) Simpson's $\frac{1}{3}$ rule is better than Simpon's $\frac{3}{8}$ rule
 - (iv) None of the above

- 2. Answer **any four** from the following questions: 2×4=8
 - (a) Show that the series $\frac{1}{1^{P}} \frac{1}{2^{P}} + \frac{1}{3^{P}} \frac{1}{4^{P}} + \dots \text{ converges for } P > 0.$
 - (b) Define bounded and unbounded sets.

 Is the set of natural numbers bounded?
 - (c) If M and N are neighbourhood of a point x, then prove that $M \cap N$ is also a neighbourhood of x.
 - (d) State Taylor's theorem with Lagrange's and Cauchy's form of remainder.
 - (e) Using Lagrange's mean value theorem prove that $|\tan^{-1} x \tan^{-1} y| \le |x y| \ \forall x, y \in \mathbb{R}$
 - (f) Solve the difference equation $u_{x+1} au_x = 0, a \ne 1$
 - (g) Write a note on numerical integration.
 - (h) Find the first three divided differences of the function $\frac{1}{x}$ for the arguments a, b, c, d.

- 3. Answer **any three** from the following questions: 5×3=15
 - (a) Show that $\lim_{n\to\infty} n^{\frac{1}{n}} = 1$
 - (b) Expand e^x by Maclaurin's infinite series.
 - (c) (i) Define limit superior and limit inferior of a bounded sequence.
 - (ii) Prove that the intersection of a finite family of open sets is open.
 - (d) Show that between any two roots of $e^x \cos x = 1$, there exists at least one root of $e^x \sin x 1 = 0$.
 - (e) State the following:
 - (i) d'Alembert's ratio text
 - (ii) Raabe's test
 - (iii) Absolute convergence of series 1

2

- (f) State and prove Simpson's $\frac{1}{3}$ rule.
- (g) Establish the relation between operator E of calculus of finite differences and differential coefficient D of differential calculus.

Also show that

 $\nabla = 1 - e^{-hD}$

where ∇ is called backward difference operation. 3+2=5

- (h) State and prove Gauss's forward interpolation formula.
- 4. Answer **any three** of the following questions: 10×3=30
 - (a) State and prove Cauchy's general principle of convergence.
 - (b) Prove that if f(x) is a function, which is
 - (i) continuous in the closed interval [a, b]

- (ii) differentiable in the open interval (a, b) and
 - (iii) f(a) = f(b), then there exists one value of x say $\xi \in]a,b[$ such that $f(\xi) = 0$

Also give the geometrical meaning of Rolle's theorem. 8+2=10

- (c) Expand log(1+x) by Maclaurin's infinite series.
- (d) (i) State Cauchy's first theorem on limits.
 - (ii) Show that

$$\lim_{n \to \infty} \left[\frac{1}{\sqrt{n^2 + 1}} + \frac{1}{\sqrt{n^2 + 2}} + \dots + \frac{1}{\sqrt{n^2 + n}} \right] = 1$$

- (iii) Define (i) monotonic sequence and (ii) derived sets.
- (e) (i) State and prove Lagrange's mean value theorem. Also give its geometrical interpretation. 5+2=7

- (ii) Show that a necessary condition for convergence of an infinite series $\sum u_n \text{ is that } \lim_{n\to\infty} \sum u_n = 0.$
- (f) State and prove Lagrange's interpolation formula for unequal intervals. Also show that the sum of the Lagrangian coefficients is unity. 7+3=10
 - (g) (i) State and prove Simpson's $\frac{3}{8}$ rule. Also state its assumptions.
 - (ii) Solve the difference equation $u_{x+1} 3^x u_x = 0$ 4
 - (h) (i) Write a note on use of various interpolation formulae. 5
 - (ii) Evaluate $\Delta^{2} \left[\frac{5x+12}{x^{2}+5x+6} \right], \text{ taking } h=1$ 5

State and prove Lagrange's mean